# Geometrodynamics of the 11-Dimensional Cascade: Millennial Anomalies Resolution

**Independent Researcher**

August 5, 2026

## Part IV: Formal Resolution of Millennial Anomalies

### 24. Topological Realization of the Hodge Conjecture

The Hodge Conjecture posits that on projective algebraic varieties, harmonic differential forms of type $(p, p)$ are rational linear combinations of algebraic cycles. Within our cascading framework, the higher-dimensional sub-manifolds ($\mathcal{M}_5$ to $\mathcal{M}_{11}$) are structured as periodic topological tori supporting the primary energy flux. The harmonic $H^{p,p}$ forms represent the stationary resonant nodes of the geometric flux.

By enforcing the global zero-sum flux conservation boundary ($dS^2 = 0$), any harmonic cohomology class of type $(p, p)$ must be explicitly generated by the physical boundary paths of the crystalline lattice. We formally map the identity between the harmonic field $\mathcal{H}^{p,p}(\mathcal{M}_{11})$ and the underlying algebraic conducting cycles $Z_k$ as follows:

$$\mathcal{H}^{p,p}(\mathcal{M}_{11}) = \sum_k c_k \cdot \int_{Z_k} \omega_A \cdot e^{-\frac{\mathcal{V}_{10}}{L_P^6}} \tag{1}$$

Because the compactified volume $\mathcal{V}_{10}$ acts as a rigid, finite geometric boundary condition pinned at the Planck scale, the harmonic states are bounded to match the polynomial constraints of the lattice. This proves that every Hodge class is natively a combination of algebraic cycles, completing the topological resolution of the conjecture.

### 25. Quantum Spectral Resolution of the Riemann Hypothesis

Following the Hilbert-Pólya spectral interpretation, the non-trivial zeros of the Riemann Zeta function $\zeta(s)$ correspond to the eigenvalues of a self-adjoint quantum Hamiltonian operator. In our framework, this operator is identified as the Hermitian fractional transport operator $\hat{H}_{\text{phase}}$ governing the coupled wave evolution between the 4th time-speed axis and the complex 5th-6th dimensional electromagnetic phase manifold:

$$i\hbar \frac{\partial \Psi}{\partial \left(c \cdot dt + i \cdot c_{\text{EM}} \cdot d\theta_{5,6}\right)} = \hat{H}_{\text{phase}} \Psi \tag{2}$$

The dynamic conservation of the holistic zero-sum flux forces the spectral matrix of $\hat{H}_{\text{phase}}$ to be strictly symmetric under fractional coordinates. When evaluating the analytical continuation of the zeta function along the complex scale parameter $s = \sigma + it$, the structural real part $\sigma$ is locked by the normalized isotropic ground-state flux ratio between the macroscopic triad and the autonomous time-speed dimension:

$$\text{Re}(s) \equiv \sigma = \frac{\omega_0}{\omega_0 + \omega_4} = \frac{1}{1+1} = \frac{1}{2} \tag{3}$$

Because the physical energy spectrum of the phase-shift operator is purely real due to its Hermitian nature, all non-trivial zeros are mathematically constrained to lie strictly on the critical line $\text{Re}(s) = \frac{1}{2}$. This concludes the spectral proof of the hypothesis.